

## Resting state functional connectivity of the basal nucleus of Meynert in humans: In comparison to the ventral striatum and the effects of age



Chiang-shan R. Li<sup>a,b,c,\*</sup>, Jaime S. Ide<sup>a,d</sup>, Sheng Zhang<sup>a</sup>, Sien Hu<sup>a</sup>, Herta H. Chao<sup>e,f</sup>, Laszlo Zaborszky<sup>g</sup>

<sup>a</sup> Department of Psychiatry, Yale University, New Haven, CT 06519, USA

<sup>b</sup> Department of Neurobiology, Yale University, New Haven, CT 06520, USA

<sup>c</sup> Interdepartmental Neuroscience Program, Yale University, New Haven, CT 06520, USA

<sup>d</sup> Department of Science and Technology, Federal University of Sao Paulo, Sao Jose dos Campos, SP 12231, Brazil

<sup>e</sup> Department of Internal Medicine, Yale University New Haven, CT 06519, USA

<sup>f</sup> Medical Service, VA Connecticut Health Care System, West Haven, CT 06516, USA

<sup>g</sup> Center for Molecular and Behavioral Neuroscience, Rutgers, NJ 07102, USA

### ARTICLE INFO

#### Article history:

Accepted 6 April 2014

Available online 13 April 2014

#### Keywords:

Functional connectivity

fMRI

Resting state

Nucleus of Meynert

Ventral striatum

Basal forebrain

### ABSTRACT

The basal nucleus of Meynert (BNM) provides the primary cholinergic inputs to the cerebral cortex. Loss of neurons in the BNM is linked to cognitive deficits in Alzheimer's disease and other degenerative conditions. Numerous animal studies described cholinergic and non-cholinergic neuronal responses in the BNM; however, work in humans has been hampered by the difficulty of defining the BNM anatomically. Here, on the basis of a previous study that delineated the BNM of post-mortem human brains in a standard stereotaxic space, we sought to examine functional connectivity of the BNM, as compared to the nucleus accumbens (or ventral striatum, VS), in a large resting state functional magnetic resonance imaging data set. The BNM and VS shared but also showed a distinct pattern of cortical and subcortical connectivity. Compared to the VS, the BNM showed stronger positive connectivity with the putamen, pallidum, thalamus, amygdala and midbrain, as well as the anterior cingulate cortex, supplementary motor area and pre-supplementary motor area, a network of brain regions that respond to salient stimuli and orchestrate motor behavior. In contrast, compared to the BNM, the VS showed stronger positive connectivity with the ventral caudate and medial orbitofrontal cortex, areas implicated in reward processing and motivated behavior. Furthermore, the BNM and VS each showed extensive negative connectivity with visual and lateral prefrontal cortices. Together, the distinct cerebral functional connectivities support the role of the BNM in arousal, saliency responses and cognitive motor control and the VS in reward related behavior. Considering the importance of BNM in age-related cognitive decline, we explored the effects of age on BNM and VS connectivities. BNM connectivity to the visual and somatomotor cortices decreases while connectivity to subcortical structures including the midbrain, thalamus, and pallidum increases with age. These findings of age-related changes of cerebral functional connectivity of the BNM may facilitate research of the neural bases of cognitive decline in health and illness.

© 2014 Elsevier Inc. All rights reserved.

### Introduction

A prominent feature of the basal forebrain is the collection of large cortically projecting neurons (basal nucleus of Meynert/BNM or magnocellular basal forebrain cell groups) that serve as the primary source of cholinergic input to the entire cortical mantle (Mufson et al., 2003; Raghanti et al., 2011; Selden et al., 1998; Wenk, 1997; Zaborszky et al., 2013, in press). In Alzheimer's and related neurodegenerative diseases, there is a severe loss of cholinergic neurons, and the decrement in cholinergic inputs to the cerebral cortex may underlie cognitive deficits that characterize these age-related conditions

(Garibotto et al., 2013). Furthermore, decades of clinical trials showed that medications enhancing cholinergic signals have efficacy in improving cognitive functions in some individuals with Alzheimer's diseases (Doody et al., 2012; Wallace and Bertrand, 2013). Despite its broad involvement in attention, memory, and other cognitive capacities, how the BNM functionally integrates with the rest of the brain is not well understood. This gap of knowledge arises at least in part from the anatomical complexity of the basal forebrain.

The BNM is interdigitated with several anatomical systems in the basal forebrain, including the ventral striopallidal system (ventral pallidum and nucleus accumbens), and cell groups underneath the globus pallidus that bridge the centromedial amygdala to the bed nucleus of the stria terminalis (so-called 'extended amygdala'; Heimer et al., 1991). This complexity has hampered precise anatomical delineation of the BNM. Recently, with cytoarchitectonics of postmortem

\* Corresponding author at: Connecticut Mental Health Center S112, 34 Park Street, New Haven CT 06519, USA. Fax: +1 203 974 7076.

E-mail address: [chiang-shan.li@yale.edu](mailto:chiang-shan.li@yale.edu) (C.R. Li).

human brains, we presented stereotaxic probabilistic maps of the basal forebrain areas containing the magnocellular cell groups (Zaborszky et al., 2008). On histological sections in ten postmortem brains, the individual compartments of the magnocellular cell groups of the basal forebrain were delineated, 3D reconstructed, and warped to a reference space in Montreal Neurological Institute (MNI) coordinates. The superposition of the cytoarchitectonic maps in the MNI brain shows intersubject variability of the various nuclei and their stereotaxic position relative to other brain structures. Both the right and left BNM showed significantly smaller volumes with increasing age (Zaborszky et al., 2008). This probabilistic map would provide a valuable tool for research of the functions of the BNM in humans that has heretofore been difficult.

Numerous studies have suggested connectivity analysis of resting state fMRI data as a useful alternative to characterizing functional architecture of a brain region. Specifically, low frequency blood oxygenation level dependent (BOLD) signal fluctuations reflect connectivity between functionally related brain regions (Biswal et al., 1995; Fair et al., 2007; Fox and Raichle, 2007). For instance, based on the findings that regions with similar functionality tend to be correlated in their spontaneous BOLD activity, investigators described functional subdivisions of many brain structures including the thalamus (Zhang et al., 2008, 2010), basal ganglia (Barnes et al., 2010), medial superior frontal cortex (Kim et al., 2010; Zhang et al., 2012), anterior cingulate cortex (Margulies et al., 2007), orbitofrontal cortex (Kahnt et al., 2012), cerebellum (O'Reilly et al., 2010), and precuneus (Cauda et al., 2010a,b; Margulies et al., 2009; Zhang and Li, 2012a).

The current study aimed to employ the probabilistic map (Zaborszky et al., 2008) and characterize whole brain functional connectivity of the BNM, as a step to understanding the systems-level functions of this basal forebrain structure in humans. In particular, we compared the functional connectivities of BNM and nucleus accumbens, a ventral striatal structure that is in proximity of the cholinergic space, in the hope of delineating the distinct roles of these two anatomically adjacent structures. In addition to their spatial proximity, both BNM and VS have been implicated in goal-directed behavior (Da Cunha et al., 2012; Grace et al., 2007; Pauli and O'Reilly, 2008; Sarter et al., 2006). In particular, evidence is accruing that the cholinergic and dopaminergic systems may interact to mediate these motivated behaviors (Lester et al., 2010; Mark et al., 2011; Threlfell and Cragg, 2011). For instance, nicotinic receptor blockade or desensitization alters neuronal bursting and dopamine outflow from the VS (Rice and Cragg, 2004). Depletion of cholinergic signals in the ventral striatum resulted in deficits in sensorimotor gating and other cognitive functions (Laplante et al., 2012). Thus, examining the shared and distinct circuits would also further our understanding of the interacting roles of BNM and VS.

Additional goals were to explore the effects of age and gender on the functional connectivities of the BNM and VS. Because the BNM is implicated in age-related cognitive changes, examining age-dependent patterns of BNM connectivity will facilitate research of the neural bases of mild cognitive impairment, Alzheimer's disease and other memory disorders.

## Materials and methods

### Resting state data

Resting-state fMRI scans were pooled from three data sets (Leiden\_2180/Leiden\_2200, Newark, and Beijing\_Zang,  $n = 144$ ), downloadable from the 1000 Functional Connectomes Project (Biswal et al., 2010), and our own data ( $n = 81$ ). In selecting the data, we tried to include as many subjects as possible in order to have more stable findings in the current study, as in our earlier work (Zhang and Li, 2014; Zhang et al., 2012). Thus, we used only datasets acquired under conditions identical to our own datasets (e.g., similar TR, all under 3 T, all eye closed). Individual subjects' images were viewed one by one to

ensure that the whole brain was covered. A total of 225 healthy subjects' resting state data (18–53 years of age; 109 men; one scan per participant; duration: 4.5–10 min) were analyzed. Table 1 summarizes these data sets.

### Imaging data preprocessing

Brain imaging data were preprocessed using Statistical Parametric Mapping (SPM 8, Wellcome Department of Imaging Neuroscience, University College London, U.K.). Images of each individual subject were first realigned (motion corrected) and corrected for slice timing. Individual structural image was normalized to an MNI (Montreal Neurological Institute) EPI (echo-planar imaging) template with affine registration followed by nonlinear transformation (Ashburner and Friston, 1999, 2005). The normalization parameters determined for the structure volume were then applied to the corresponding functional image volumes for each subject. Finally, the images were smoothed with a Gaussian kernel of 8 mm at Full Width at Half Maximum.

Additional preprocessing was applied to reduce spurious BOLD variances that were unlikely to reflect neuronal activity (Fair et al., 2007; Fox and Raichle, 2007; Fox et al., 2005; Rombouts et al., 2003). The sources of spurious variance were removed through linear regression by including the signal from the ventricular system, white matter, and whole brain, in addition to the six parameters obtained by rigid body head motion correction. First-order derivatives of the whole brain, ventricular and white matter signals were also included in the regression.

Cordes and colleagues suggested that BOLD fluctuations below a frequency of 0.1 Hz contribute to regionally specific BOLD correlations (Cordes et al., 2001). Thus, we applied a temporal band-pass filter ( $0.009 \text{ Hz} < f < 0.08 \text{ Hz}$ ) to the time course in order to obtain low-frequency fluctuations, as in previous studies (Fair et al., 2007; Fox and Raichle, 2007; Fox et al., 2005; Lowe et al., 1998).

### Head motion

As extensively investigated in Van Dijk et al., 2012, microhead motion ( $>0.1 \text{ mm}$ ) is an important source of spurious correlations in resting state functional connectivity analysis. Therefore, we applied a “scrubbing” method proposed by Power et al. (2012) and successfully applied in previous studies (Power et al., 2012; Smyser et al., 2010; Tomasi and Volkow, 2014) to remove time points affected by head motions. Briefly, for every time point  $t$ , we computed the *framewise displacement* given by  $FD(t) = |\Delta d_x(t)| + |\Delta d_y(t)| + |\Delta d_z(t)| + r|\alpha(t)| + r|\beta(t)| + r|\gamma(t)|$ , where  $(d_x, d_y, d_z)$  and  $(\alpha, \beta, \gamma)$  are the translational and rotational movements, respectively, and  $r (= 50 \text{ mm})$  is a constant that approximates the mean distance between center of MNI space and the cortex and transform rotations into displacements (Power et al., 2012). The second head movement metric was the root mean square variance (DVARs) of

**Table 1**

Demographic information and imaging parameters of the resting-state functional MRI data obtained from the image repository for the 1000 Functional Connectomes Project and our laboratory.

Dataset	Subjects	Ages (years)	Time points	TR (s)	Slice acquisition order
Beijing_Zang	31 M/66 F	18–26	225	2	Interleaved ascending
Leiden_2180	10 M/0 F	20–27	215	2.18	Sequential descending
Leiden_2200	11 M/8 F	18–28	215	2.2	Sequential descending
Newark	9 M/9 F	21–39	135	2	Interleaved ascending
Our own	48 M/33 F	19–53	295	2	Interleaved ascending

Note: M, males; F, females; and TR: repetition time.

the differences in % BOLD intensity  $I(t)$  between consecutive time points across brain voxels, computed as follows:  $DVARF(t) = \sqrt{|I(t) - I(t-1)|^2}$ , where the brackets indicate the mean across brain voxels. Finally, to compute each subject's correlation map, we removed every time point that exceeded the head motion limit  $FD(t) > 0.5$  mm or  $DVARF(t) > 0.5\%$  (Power et al., 2012; Tomasi and Volkow, 2014). On average, 1% of the time points were removed across subjects, and 2 (out of 225) individuals were excluded from further analyses because of excessive head movements.

#### Seed regions: basal nucleus of Meynert and ventral striatum

A mask of the basal nucleus of Meynert was created based on a stereotaxic probabilistic map of magnocellular cell groups in the basal forebrain (Zaborszky et al., 2008). Briefly, a T1-weighted MRI scan of  $1.17 \times 1 \times 1$  mm was obtained of each individual brain ( $n = 10$ ) before histological processing. The outlines of various basal forebrain compartments were traced on 2D images of silver-stained (Merker, 1983) histological sections ( $20 \mu\text{m}$  thick, 1.2 mm apart) with a final resolution of  $7000 \times 6000$  pixels. The outlines were processed as contour line for each histological section. As described in Zaborszky et al., 2008, we used a modified version of the Ch1–Ch4 nomenclature of Mesulam et al. (1983) to delineate the magnocellular basal forebrain cell groups. Ch1–Ch2 compartment largely corresponds to the medial septum and the diagonal band of Broca. The Ch3 group corresponds to the horizontal nucleus of the diagonal band of Broca as defined by Hedreen et al. (1984). Cell aggregates in the subcommissural–sublenticular region largely correspond to the basal nucleus of Meynert as defined by Vogels et al. (1990) and De Lacalle et al. (1991) and were collectively defined as “Ch4”. Subdivisions of the “Ch4” as suggested by Mesulam et al. (1983) were not delineated.

Magnocellular neurons in the basal forebrain often constitute cell aggregates or clusters that facilitate their recognition. As a rule, borders of delineated areas were drawn where the number of large cells decreased precipitously. Details of reconstruction of these nuclei on a 3D template were described in Zaborszky et al., 2008. Briefly, a 3D volume was reconstructed, with a known algorithm from the digitized histological sections using the MRI scans of the fixed brain as a shape reference with a resolution of 1 mm isotropic (Hoemke, 2006; Schormann and Zilles, 1998). Each 3D reconstructed histological volume was spatially normalized to the single subject T1-weighted MNI reference brain (Collins et al., 1994; Holmes et al., 1998) using a


combination of affine linear and non-linear, elastic transformations (Henn et al., 1997; Hoemke, 2006). Based on the 3D whole-brain reconstructions, volume and shape of the magnocellular cell groups were computed. Probability maps were computed separately for each Ch compartment. In this study we used the mask that corresponds to the Ch4 cell groups. These probabilistic maps describe the relative frequency with which the same basal forebrain area (e.g., Ch4) was represented in each voxel of the reference space.

In primates, the nucleus accumbens merges imperceptibly with the rostroventral parts of the caudate nucleus and putamen. We used both cytoarchitectonic and topographical criteria to generate a ventral striatal mask, that largely corresponds to the nucleus accumbens using the following criteria for delineation on enlarged images of the histological sections: there is a higher cell density in the nucleus accumbens than in dorsal striatum; the slightly smaller cells are of round/oval shape; and very few or no fiber bundles pass through the area. Caudally, the lower edge of the ventral pallidum, which appears dark due to its high iron content in silver-stained sections, served as the dorsal border. However, ventral pallidal islands are invaginated into the nucleus accumbens; therefore, caudally the ventral striatal volume inevitably contains some ventral pallidal tissue. In the region of the anterior perforated substance, the nucleus accumbens extends almost to the ventral surface of the brain. Cell groups such as the ‘interface islands’ (Heimer, 2000) were excluded, provided that they are surrounded by thin glial layer.

These two regions of interest are shown in Fig. 1. We excluded from each region the overlapping voxels (yellow) to form the BNM and VS seed regions for connectivity analyses.

#### Orthogonalization

To highlight the distinct functional connectivity patterns between basal nucleus of Meynert (BNM) and the ventral striatum (VS), we constructed BNM and VS seed masks by excluding the overlapping area between them as depicted in Fig. 1. Additionally, the Gram–Schmidt method was used to orthogonalize the BNM and VS signals (Di Martino et al., 2008; Margulies et al., 2007; Tomasi and Volkow, 2014). The orthogonalization process highlights the variances unique to BNM and VS signals, by reducing the common confounding fluctuations. However, when querying cerebral functional connectivity of a region of interest (ROI), investigators are primarily interested in connectivities of the ROI without considering whether these results are



**Fig. 1.** Seed regions: basal nucleus of Meynert (red) and ventral striatum (green), with overlapping voxels in yellow, shown on axial (from  $z = -26$  to  $+2$ ) and coronal (from  $y = -10$  to  $+26$ ) sections. Overlapping voxels were removed from each seed region prior to whole brain regressions.

independent of those of other brain regions. Therefore, we presented the results obtained with the original time series in the main text and those with the orthogonalized time series in the Supplementary data. In practice, orthogonalization is expected to have little impact when comparing the BNM and VS correlation maps, since common variations are eliminated in the statistical test.

#### *Seed region-based linear correlation and random effects analysis*

The BOLD time courses were averaged spatially over each of the two seeds. For individual subjects, we computed the correlation coefficient between the averaged time course of each seed region and the time courses of all other brain voxels. To assess and compare the resting state “correlograms,” we converted these image maps, which were not normally distributed, to z score maps by Fisher’s z transform (Berry and Mielke, 2000; Jenkins and Watts, 1968):  $z = 0.5 \log_e[(1 + r)/(1 - r)]$ . The Z maps were used in group random effect analyses. We performed one-sample *t* test each on the Z maps of BNM and VS and paired-sample *t* test comparing the two Z maps.

#### *Age dependent changes and gender differences in BNM and VS connectivity*

We performed a simple regression of the Z maps against age, each for the BNM and VS, to identify age-related changes of functional connectivity in the two structures. To examine gender differences, we compared men and women with age as a covariate in an analysis of variance, each for the BNM and VS. All results were reported for a corrected threshold.

## Results

### *Whole brain functional connectivity of the nucleus of Meynert and ventral striatum*

For each seed region, we performed one sample *t*-test of the Z maps (positively and negatively correlated regions) across the group ( $n = 223$ ). Regions functionally connected with nucleus of Meynert (BNM) and ventral striatum (VS) are presented in Figs. 2 and 3, respectively.

The BNM showed positive connectivity with the medial prefrontal cortex, including dorsal/rostral/perigenual/subgenual anterior cingulate cortex (ACC), supplementary motor area (SMA) as well as pre-SMA, medial orbitofrontal cortex, inferior temporal pole, hippocampus, amygdala, insula, thalamus, midbrain, and basal ganglia. The BNM showed negative connectivity with the cuneus, parahippocampal gyri, precuneus, somatosensory cortex, posterior parietal cortices, middle and lateral frontal cortices, including the dorsolateral prefrontal cortex and frontal eye field, and cerebellum (Fig. 2).

The VS showed positive connectivity with the medial orbitofrontal cortex, rectus gyrus, subcallosal gyrus, inferior temporal pole, amygdala, hippocampus, perigenual/subgenual ACC, and ventral part of the caudate head. The VS showed negative connectivity with the visual cortices, posterior parietal cortex, temporoparietal junction, superior temporal gyri, precentral cortex, lateral frontal cortices including the dorsolateral prefrontal cortex, inferior frontal cortices, posterior thalamus/pulvinar, parahippocampal gyrus, and cerebellum (Fig. 3). These results are consistent with previous findings of resting state VS connectivity with the medial orbitofrontal cortex and ventral caudate (Baliki et al., 2013; Cauda et al., 2011; Di Martino et al., 2008; Gopinath et al., 2011), amygdala and midbrain (Cauda et al., 2011), and hippocampus (Cauda et al., 2011; Kahn and Shohamy, 2013).

We performed a paired *t*-test to compare functional connectivity of the BNM and VS (Fig. 4). Compared to the VS, the BNM showed greater connectivity with the medial frontal cortex, including rostral/dorsal ACC, SMA, pre-SMA, dorsolateral prefrontal cortex, inferior frontal cortex, lateral orbitofrontal cortex, inferior parietal lobule, temporo-

parietal junction, insula, thalamus, dorsal caudate, midbrain, and cerebellum. Compared to the BNM, VS showed greater connectivity with the medial orbitofrontal cortex, ventral caudate, subcallosal/rectus gyri, olfactory sulcus, cuneus, precuneus, posterior cingulate cortex, parahippocampal gyrus, middle/inferior occipital cortex, and postcentral gyri. We summarize these results in Tables 2 and 3, where we further distinguish whether a contrast arises from differences in positive or negative connectivity.

#### *The effects of age and gender on BNM and VS connectivity*

In a simple regression, BNM connectivity to the midbrain in the area of the ventral tegmental area and substantia nigra (MNI coordinates  $[-9, -16, -14]$ ,  $Z = 5.41$ ; and  $[6, -19, -20]$ ,  $Z = 5.08$ ,  $1215 \text{ mm}^3$ ), left pallidum and parahippocampal gyrus ( $[-30, -7, -11]$ ,  $Z = 5.62$ ,  $1026 \text{ mm}^3$ ), and right pallidum ( $[27, -7, -5]$ ,  $Z = 5.04$ ,  $189 \text{ mm}^3$ ) showed a positive correlation with age (Fig. 5). In contrast, BNM connectivity to the bilateral motor cortices ( $[-33, -37, 70]$ ,  $Z = 5.63$ ; and  $[-45, -16, 61]$ ,  $Z = 4.85$ ,  $2187 \text{ mm}^3$ ; and  $[45, -25, 64]$ ,  $Z = 5.44$ ,  $1161 \text{ mm}^3$ ) and right visual cortex ( $[15, -94, 25]$ ,  $Z = 5.59$ ,  $2133 \text{ mm}^3$ ) showed a negative correlation with age. At the same, corrected threshold, VS connectivities did not show any age dependence.

In a covariance analysis with age as a covariate, we examined gender differences. Men and women did not differ in BNM or VS connectivity at the same, corrected threshold.

## Discussion


### *Functional connectivity of the nucleus of Meynert and ventral striatum*

Compared to the VS, the BNM showed stronger positive connectivity with the putamen, pallidum, thalamus, amygdala and midbrain, cerebellum, as well as the anterior cingulate cortex, supplementary motor area and pre-supplementary motor area, a network of brain regions that respond to salient stimuli and orchestrate goal-directed behavior. In contrast, the VS showed stronger positive connectivity with the ventral caudate and medial orbitofrontal cortex, areas implicated in reward processing, compared to the BNM. Together, the distinct cerebral functional connectivities support the role of the BNM in arousal and cognitive motor control and the VS in reward related behavior.

The BNM shows connectivity with a selective set of cortical and subcortical structures, consistent with anatomical studies in rodents and non-human primates (Chandler and Waterhouse, 2012; Mesulam et al., 1984; Zaborszky et al., in press). In particular, studies in monkeys showed that the density of cholinergic fibers and choline acetyltransferase activities are particularly rich in agranular areas of the insula, caudal orbitofrontal cortex (OFC), temporal pole, and parahippocampal region (Mesulam et al., 1984, 1986), a pattern mirrored by the current findings of regions positively connected to the BNM (Fig. 2). The VS receives extensive projections from the OFC (Haber et al., 2006), amygdala, and dysgranular part of the insula that process taste, olfaction and other visceral information (Fudge et al., 2005; Nakano et al., 1999), in accord with VS responses to primary reinforcers.

### *Functional implications for the basal nucleus of Meynert and ventral striatum*

The BNM, along with the pulvinar and amygdala, responds to salient stimuli (Morris et al., 1997). Manipulations of cholinergic signaling alter neural responses to salient stimuli in a variety of behavioral paradigms, including odd-ball tasks (Klinkenberg et al., 2013). For instance, during auditory conditioning in humans, augmenting cholinergic signaling with physostigmine (an inhibitor of acetylcholinesterase) enhances processing of behaviorally irrelevant stimuli and attenuates differential conditioning-related cortical activations (Thiel et al., 2002).




**Fig. 2.** Brain areas that show positive (warm color) and negative (cool color) functional connectivity to the basal nucleus of Meynert; one-sample  $t$  test,  $p < 0.05$ , corrected for familywise error of multiple comparisons. White matter voxels are excluded with a mask.

Scopolamine, an antagonist of muscarinic acetylcholine receptors, impairs categorization of new but not learnt visual stimuli in monkeys (Aggelopoulos et al., 2011). Donepezil, a cholinesterase inhibitor, facilitates reaction time to targets following a cue that instructs voluntary shift of attention (Rokem et al., 2010). In rodents, increasing cholinergic activity in gustatory cortex enhances the salience of a familiar, conditioned stimulus in taste aversion learning (Clark and Bernstein, 2009). Cholinergic deafferentation lowers the signal-to-noise ratio of target evoked responses in posterior parietal neurons (Broussard et al.,

2009). Infusion of a cholinergic immunotoxin in the basal forebrain compromises cued target detection (Bushnell et al., 1998). A study using microdialysis demonstrated that acetylcholine release in the frontal cortex and hippocampus is elicited by behavioral conditioning to salient stimuli (Acquas et al., 1996). Taken together, numerous studies support a role of the BNM in processing salient and novel stimuli for learning and cognitive motor control (Wenk, 1997).


Although earlier studies largely focused on the role of the ventral striatum (VS) in reward processing, a growing body of work has



**Fig. 3.** Brain areas that show positive (warm color) and negative (cool color) functional connectivity to the ventral striatum; one-sample *t* test,  $p < 0.05$ , corrected for familywise error of multiple comparisons. White matter voxels are excluded with a mask.

suggested that the VS responds not only to rewarding but more generally to stimuli that are just salient (Jensen et al., 2003, 2007; Litt et al., 2011; Pohlack et al., 2012; Veldhuizen et al., 2011; Zink et al., 2003, 2006; see also Brooks and Berns, 2013 for a review). For instance, the VS responds to painful stimulation as well as anticipation of such aversive stimuli (Jensen et al., 2003; Zink et al., 2006). A critical issue then is to distinguish functional roles of the BNM and VS in saliency processing.

The current findings on the differences in functional connectivity may help address this question. Compared to the VS, the BNM shows stronger positive connectivity with the medial frontal and fronto-parietal cortices, as well as the thalamus, insula, basal ganglia, and midbrain – areas that form a task-related network (Robinson et al., 2009; Zhang et al., 2008), as well as stronger negative connectivity with the precuneus, a core area of the default mode network (Raichle and



**Fig. 4.** Brain areas that show differences in functional connectivity to the basal nucleus of Meynert (BNM) versus ventral striatum (VS): BNM > VS (warm color); VS > BNM (cool color); paired-sample *t* test,  $p < 0.05$ , corrected for familywise error of multiple comparisons.

Snyder, 2007; Raichle et al., 2001; Zhang and Li, 2012a). These differences suggest that, compared to the VS, the BNM is more directly and readily engaged in task related processing.

This difference in functional connectivity is consistent with a general role of the BNM in task-related behavior such as perception and perceptual motor learning and memory (Chen et al., 2012; Conner et al., 2010; Froemke et al., 2013; Goard and Dan, 2009; Leach et al., 2013; Miasnikov

and Weinberger, 2012). In particular, the BNM may mediate learning and memory even when such processes are motivationally neutral. In behavioral conditioning of rats, specific associative memory can be induced by direct activation of the BNM without detectable motivational effects as in seeking reward or avoiding punishment (Miasnikov et al., 2008). Indeed, neurons in the BNM did not respond to primary reinforcers, positive (juice) or negative (salt water), in monkeys (Wilson

**Table 2**Regions showing greater connectivity with basal nucleus of Meynert (BNM) as compared to ventral striatum (VS); paired *t*-test, *n* = 223.

Volume (mm <sup>3</sup> )	Peak voxel (Z)	MNI coordinate			Side	Identified brain region	Connectivity	
		x	y	z			BNM	VS
365472	inf	24	-1	-8	R	Putamen/Pallidum/Thalamus/Midbrain/Cerebellum/Amygdala/Insula/TP/IOFC	++	+/-/-
	inf	-24	-4	-8	L	Putamen/Pallidum/Thalamus/Midbrain/Cerebellum/Amygdala/Insula/TP/IOFC	++	+/-/-
	inf	9	11	34	R	ACC/SMA/pre-SMA	+	~/-
	inf	-9	14	37	L	ACC/SMA/pre-SMA	+	~/-
4563	7.48	63	-31	28	R	Temporo-parietal junction/Supramarginal G	+/-	-/-
	7.33	-66	-28	25	L	Temporo-parietal junction/Supramarginal G	+/-	-/-
1026	6.60	63	-46	1	R	Middle temporal G/Inferior frontal G/DLPFC	+	-
729	6.05	-60	-55	1	L	Middle temporal G/Inferior frontal G/DLPFC	+	-

Note: inf: infinity; R: right; L: left; TP: temporal pole; IOFC: lateral orbitofrontal cortex; G: gyrus; ACC: anterior cingulate cortex; SMA: supplementary motor area; DLPFC: dorsolateral prefrontal cortex; +/- indicates positive and negative connectivity (with + +/- - for stronger connectivity), and - indicates no significant connectivity at one-sample *t* test, *p* < 0.05, FWE corrected.

and Rolls, 1990). While the BNM responds to visual or acoustic stimuli that signal reinforcers, it does not respond to the reinforcers themselves (Wilson and Ma, 2004). Other work suggests that the BNM neurons encode saliency irrespective of valence of the stimuli (Lin and Nicolelis, 2008). These results are concordant with a learning-related or memory-promoting role for the BNM that places it “downstream” of motivational systems (Miasnikov et al., 2008), consistent with the anatomical finding that, the cholinergic neurons in basal forebrain receive input from the nucleus accumbens (Zaborszky and Cullinan, 1992). Through its extensive cortical projections, the BNM may mediate attentional effort in a circuitry that integrates brain systems involved in modulation of input functions, incentive processing and performance monitoring (Sarter et al., 2006).

On the other hand, one is cautioned not to over-interpret the differences of BNM and VS in saliency- and reward-related processing. For instance, reward is intrinsically salient, and cerebral structures such as the medial orbitofrontal cortex, with positive connectivity to both the BNM and VS, is involved in processing reward and emotionally salient stimuli (Hardin et al., 2009; Kühn and Gallinat, 2012; Liu et al., 2011; Rothkirch et al., 2012). Likewise, as discussed earlier, although VS is predominantly implicated in reward processing, there is accumulating evidence that VS responds to salient, non-rewarding stimuli (Esslinger et al., 2013; Jensen et al., 2007; Pohlack et al., 2012; Spicer et al., 2007; Zink et al., 2003, 2006). Future research may take advantage of the current findings in further delineating the respective roles of BNM and VS in motivated behavior.

#### Age-related connectivity of the nucleus of Meynert

Previous imaging studies suggest that aging is related in a complicated way to cerebral functional connectivities, with different brain areas increasing and decreasing with age in connectivity to regions of interest (Bernard et al., 2013; Campbell et al., 2013; Hafkemeijer et al., 2013; Hoffstaedter et al., 2014; Liu et al., 2012; Roski et al., 2013; Taniwaki et al., 2007; Ystad et al., 2010; see also Ferreira and Busatto, 2013 for a review). Here, we observed that age is associated with increased BNM connectivity to subcortical structures including the midbrain and

pallidum and decreased connectivity to somatomotor and visual cortices. Although statistically highly significant, these results are not straightforward to interpret without a functional concomitant. Given that the current BOLD time series were obtained during resting state, one is tempted to speculate that these changes are not related to specific contextual or task conditions. Future studies are needed to examine the relationship between this contrasting pattern of increased and decreased connectivity to subcortical and cortical structures and cognitive motor performance (Hu et al., 2012, 2013) and evaluate whether and to what extent these changes may reflect a compensatory process.

In over 1000 adolescents and young adults, Tomasi and Volkow characterized age-related increases in the functional connectivity of the ventral tegmental area (VTA) with limbic regions and with the default mode network and decreases in the connectivity of the substantia nigra (SN) with motor and medial temporal cortices (Tomasi and Volkow, 2014). In terms of mean strength and statistical significance of connectivity, the VTA demonstrated bilateral positive connectivity with the nucleus accumbens, hippocampus, parahippocampus, globus pallidus, caudate, cerebellar vermis, and anterior insula, as well as the angular gyrus, inferior frontal cortex, and anterior cingulate cortex, and negative connectivity with the inferior occipital cortex. The SN demonstrated bilateral positive connectivity with the globus pallidus, subthalamic nucleus, thalamus, and cerebellar vermis, as well as the Broca's and Wernicke's areas, anterior cingulate cortex, and supplementary motor area, and negative connectivity with orbitofrontal, temporal, and occipital cortices, ventral precuneus, and angular gyrus. Thus, there is much overlap between subcortical and cortical connectivities between the VTA/SN and BNM/VS, suggesting the need in future studies to investigate early developmental changes in functional connectivity of the BNM/VS and their relevance to clinical conditions that manifest early during development.


#### Potential clinical implications

In addition to implications for our understanding of the cognitive deficits in Alzheimer's disease and other forms of age-related cognitive decline, the current findings may also facilitate research of the

**Table 3**Regions showing greater connectivity with ventral striatum (VS) as compared to the basal nucleus of Meynert (BNM); paired *t*-test, *n* = 223.

Volume mm <sup>3</sup>	Peak voxel Z	MNI coordinate			Side	Identified brain region	Connectivity	
		x	y	z			BNM	VS
326160	inf	12	17	-11	R	ventral Caudate/mOFC/Subcallosal G/Rectus G/Olfactory S	~/-	++
	inf	-9	14	-14	L	ventral Caudate/mOFC/Subcallosal G/Rectus G/Olfactory S	~/-	++
	inf	-18	-49	22	L	pCG/Cuneus/Precuneus/Parahippocampal G	--	~/-
	inf	18	-46	25	R	pCG/Cuneus/Precuneus/Parahippocampal G	--	~/-
1512	6.47	42	-28	64	R	Postcentral G	-	~
864	5.91	-39	-31	67	L	Postcentral G	-	~

Note: inf: infinity; R: right; L: left; mOFC: medial orbitofrontal cortex; G: gyrus; S: sulcus; pCG: posterior cingulate gyrus; +/- indicates positive and negative connectivity (with + +/- - indicating stronger connectivity), and - indicates no significant connectivity at one-sample *t* test, *p* < 0.05 FWE corrected.



**Fig. 5.** Brain areas with functional connectivity to the basal nucleus of Meynert (BNM) that correlates positively (warm color) and negatively (cool color) with age.  $p < 0.05$ , corrected for familywise error of multiple comparisons.

cholinergic mechanisms of drug and alcohol addiction (Adinoff et al., 2010; Kurosawa et al., 2013; Miki et al., 2014; Paolone et al., 2013; Williams and Adinoff, 2008). Altered cholinergic neurotransmission is known to play a critical role in nicotine addiction (Dani and Balfour, 2011). More broadly, nicotinic acetylcholine receptors are heavily expressed in limbic dopamine circuitry and the medial habenula–interpeduncular nucleus complex, which are critical mediators of reinforced and addictive behavior (Leslie et al., 2013; Mark et al., 2011).

For instance, behavioral performance that requires sustained attention is associated with extracellular levels of cortical acetylcholine. Rats that are prone to attribute incentive salience to reward cues (“sign trackers”), relative to “goal-trackers”, are vulnerable to developing obesity and addiction (Paolone et al., 2013). Because of its interaction of the limbic reward and prefrontal cognitive control circuits, cholinergic neurotransmission has been a target for pharmacotherapy development for stimulant addiction (Sofuoglu and Mooney, 2009). Thus, the current

findings of the systems level functional connectivity of the BNM will help future studies to pursue the etiology and treatment of addiction on these related fronts.

Furthermore, the VS and its cerebral connectivity has been implicated in various neuropsychiatric disorders such as autism (Delmonte et al., 2013; Di Martino et al., 2011) or psychological processes of clinical interest such as consciousness (Cauda et al., 2009a) and pain (Baliki et al., 2013; Cauda et al., 2009b, 2010a,b). For instance, with differential functional and structural connectivities, the shell and core regions of the VS may each mediate prediction of monetary reward and cessation of pain (Baliki et al., 2013). Functional connectivity of the VS and hippocampus may allow integration of motivational significance and memory to guide behavior (Kahn and Shohamy, 2013), a process of relevance to depression and psychomotor retardation. Notably, animal studies have also implicated the BNM in pain (Zhang et al., 2010) and anesthesia (Laalou et al., 2008); thus, the current findings may shed light on future studies to directly contrast the roles of VS and BNM in these neural processes of clinical significance.

#### A methodological consideration

Negative functional connectivity has been observed and reported since the very beginning of the resting state fMRI study (Biswal et al., 1995). Negative functional connectivity, also called anti-correlation, represents negative cross-correlation in spontaneous BOLD signal between two brain regions. It was suggested that the global signal regression, a common step of data preprocessing in seed region based functional connectivity analyses, is a likely cause of anti-correlated functional networks (Murphy et al., 2009; Weissenbacher et al., 2009). However, recent investigations demonstrated that the negative correlations are not an artifact but have biological origins (Chai et al., 2012; Chen et al., 2011; Fox et al., 2009). For instance, negative functional connectivity is associated predominantly with long range connections and correlates with the shortest path length in the human brain network (Chen et al., 2011; Scholvinck et al., 2010; Schwarz and McGonigle, 2011). Indeed, the negative correlations between brain regions with presumably opposing functional roles have been consistently observed in different studies (Chen et al., 2011; Fox et al., 2005; Fransson, 2005; Greicius et al., 2003; Kelly et al., 2008; Uddin et al., 2009), including those using independent component analysis, which does not involve global signal regression (Cole et al., 2010; Zhang and Li, 2012b; Zuo et al., 2010). Furthermore, the existence of the negative functional connectivity was also suggested by computational simulations of cerebral network activities in both monkeys and humans (Deco et al., 2009; Honey et al., 2007; Izhikevich and Edelman, 2008) and supported by simultaneous recording of unit activity and local field potential from task-positive and task-negative (default mode) networks in cats (Popa et al., 2009). Together, these earlier studies suggest functional significance of negative functional connectivity.

#### Acknowledgments

This study was supported by NIH grants R01DA023248, K02DA026990, R21AA018004, and R01AA021449 to C-S.R.L. and NINDS grant 23945 to LZ. The NIH had no further role in the study design; in the collection, analysis and interpretation of data; in the writing of the report; or in the decision to submit the paper for publication. J.S.I. was supported by Young Researcher Award (Sao Paulo Research Foundation FAPESP 2011/08573-4). We thank investigators of the 1000 Functional Connectomes Project and those who shared the data set for making this study possible.

#### Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.neuroimage.2014.04.019>.

#### References

- Acqas, E., Wilson, C., Fibiger, H.C., 1996. Conditioned and unconditioned stimuli increase frontal cortical and hippocampal acetylcholine release: effects of novelty, habituation, and fear. *J. Neurosci.* 16 (9), 3089–3096 (May 1).
- Adinoff, B., Devous Sr., M.D., Williams, M.J., Best, S.E., Harris, T.S., Minhajuddin, A., Zielinski, T., Cullum, M., 2010. Altered neural cholinergic receptor systems in cocaine-addicted subjects. *Neuropsychopharmacology.* 35 (7), 1485–1499 (Jun).
- Aggelopoulos, N.C., Liebe, S., Logothetis, N.K., Rainer, G., 2011. Cholinergic control of visual categorization in macaques. *Front. Behav. Neurosci.* 5, 73 (Nov 15).
- Ashburner, J., Friston, K.J., 1999. Nonlinear spatial normalization using basis functions. *Hum. Brain Mapp.* 7, 254–266.
- Ashburner, J., Friston, K.J., 2005. Unified segmentation. *Neuroimage* 26, 839–851.
- Baliki, M.N., Mansour, A., Baria, A.T., Huang, L., Berger, S.E., Fields, H.L., Apkarian, A.V., 2013. Parceling human accumbens into putative core and shell dissociates encoding of values for reward and pain. *J. Neurosci.* 33 (41), 16383–16393 (Oct 9).
- Barnes, K.A., Cohen, A.L., Power, J.D., Nelson, S.M., Dosenbach, Y.B., Miezin, F.M., Petersen, S.E., Schlaggar, B.L., 2010. Identifying basal ganglia divisions in individuals using resting-state functional connectivity MRI. *Front. Syst. Neurosci.* 4, 18.
- Bernard, J.A., Peltier, S.J., Wiggins, J.L., Jaeggi, S.M., Buschkuhl, M., Fling, B.W., Kwak, Y., Jonides, J., Monk, C.S., Seidler, R.D., 2013. Disrupted cortico-cerebellar connectivity in older adults. *Neuroimage.* 83, 103–119 (Dec).
- Berry, K.J., Mielke Jr., P.W., 2000. A Monte Carlo investigation of the Fisher Z transformation for normal and nonnormal distributions. *Psychol. Rep.* 87, 1101–1114.
- Biswal, B., Yetkin, F.Z., Haughton, V.M., Hyde, J.S., 1995. Functional connectivity in the motor cortex of resting human brain using echo-planar MRI. *Magn. Reson. Med.* 34, 537–541.
- Biswal, B.B., Mennes, M., Zuo, X.N., Gohel, S., Kelly, C., Smith, S.M., Beckmann, C.F., Adelstein, J.S., Buckner, R.L., Colcombe, S., Dogonowski, A.M., Ernst, M., Fair, D., Hampson, M., Hoptman, M.J., Hyde, J.S., Kiviniemi, V.J., Kotter, R., Li, S.J., Lin, C.P., Lowe, M.J., Mackay, C., Madden, D.J., Madsen, K.H., Margulies, D.S., Mayberg, H.S., McMahon, K., Monk, C.S., Mostofsky, S.H., Nagel, B.J., Pekar, J.J., Peltier, S.J., Petersen, S.E., Riedel, V., Rombouts, S.A., Rypma, B., Schlaggar, B.L., Schmidt, S., Seidler, R.D., Siegle, G.J., Sorg, C., Teng, G.J., Veijola, J., Villringer, A., Walter, M., Wang, L., Weng, X.C., Whitfield-Gabrieli, S., Williamson, P., Windischberger, C., Zang, Y.F., Zhang, H.Y., Castellanos, F.X., Milham, M.P., 2010. Toward discovery science of human brain function. *Proc. Natl. Acad. Sci. U. S. A.* 107, 4734–4739.
- Brooks, A.M., Berns, G.S., 2013. Aversive stimuli and loss in the mesocorticolimbic dopamine system. *Trends Cogn. Sci.* 17 (6), 281–286 (Jun).
- Broussard, J., Karelna, K., Sarter, M., Givens, B., 2009. Cholinergic optimization of cue-evoked parietal activity during challenged attentional performance. *Eur. J. Neurosci.* 29 (8), 1711–1722 (Apr).
- Bushnell, P.J., Chiba, A.A., Oshiro, W.M., 1998. Effects of unilateral removal of basal forebrain cholinergic neurons on cued target detection in rats. *Behav. Brain Res.* 90 (1), 57–71 (Jan).
- Campbell, K.L., Grigg, O., Saverino, C., Churchill, N., Grady, C.L., 2013. Age differences in the intrinsic functional connectivity of default network subsystems. *Front. Aging Neurosci.* 5, 73 (Nov 14).
- Cauda, F., Micon, B.M., Sacco, K., Duca, S., D'Agata, F., Geminiani, G., Canavero, S., 2009a. Disrupted intrinsic functional connectivity in the vegetative state. *J. Neurol. Neurosurg. Psychiatry* 80 (4), 429–431 (Apr).
- Cauda, F., Sacco, K., Duca, S., Cocito, D., D'Agata, F., Geminiani, G.C., Canavero, S., 2009b. Altered resting state in diabetic neuropathic pain. *PLoS One* 4 (2), e4542.
- Cauda, F., D'Agata, F., Sacco, K., Duca, S., Cocito, D., Paolasso, I., Isoardo, G., Geminiani, G., 2010a. Altered resting state attentional networks in diabetic neuropathic pain. *J. Neurol. Neurosurg. Psychiatry* 81 (7), 806–811 (Jul).
- Cauda, F., Geminiani, G., D'Agata, F., Sacco, K., Duca, S., Bagshaw, A.P., Cavanna, A.E., 2010b. Functional connectivity of the posteromedial cortex. *PLoS One* 5 (9), e13030 (Sep 30).
- Cauda, F., Cavanna, A.E., D'Agata, F., Sacco, K., Duca, S., Geminiani, G.C., 2011. Functional connectivity and coactivation of the nucleus accumbens: a combined functional connectivity and structure-based meta-analysis. *J. Cogn. Neurosci.* 23 (10), 2864–2877 (Oct).
- Chai, X.J., Castanon, A.N., Ongur, D., Whitfield-Gabrieli, S., 2012. Anticorrelations in resting state networks without global signal regression. *Neuroimage* 59, 1420–1428.
- Chandler, D., Waterhouse, B.D., 2012. Evidence for broad versus segregated projections from cholinergic and noradrenergic nuclei to functionally and anatomically discrete subregions of prefrontal cortex. *Front. Behav. Neurosci.* 6, 20 (May 21).
- Chen, G., Chen, G., Xie, C., Li, S.J., 2011. Negative functional connectivity and its dependence on the shortest path length of positive network in the resting-state human brain. *Brain Connect.* 1, 195–206.
- Chen, N., Sugihara, H., Sharma, J., Perea, G., Petracvic, J., Le, C., Sur, M., 2012. Nucleus basalis-enabled stimulus-specific plasticity in the visual cortex is mediated by astrocytes. *Proc. Natl. Acad. Sci. U. S. A.* 109 (41), E2832–E2841 (Oct 9).
- Clark, E.W., Bernstein, I.L., 2009. Boosting cholinergic activity in gustatory cortex enhances the salience of a familiar conditioned stimulus in taste aversion learning. *Behav. Neurosci.* 123 (4), 764–771 (Aug).
- Cole, D.M., Beckmann, C.F., Long, C.J., Matthews, P.M., Durcan, M.J., Beaver, J.D., 2010. Nicotine replacement in abstinent smokers improves cognitive withdrawal symptoms with modulation of resting brain network dynamics. *Neuroimage* 52, 590–599.
- Collins, D.L., Neelin, P., Peters, T.M., Evans, A.C., 1994. Automatic 3D intersubject registration of MR volumetric data in standardized Talairach space. *J. Comput. Assist. Tomogr.* 18, 192–205.
- Conner, J.M., Kulczycki, M., Tuszynski, M.H., 2010. Unique contributions of distinct cholinergic projections to motor cortical plasticity and learning. *Cereb. Cortex* 20 (11), 2739–2748 (Nov).

- Cordes, D., Haughton, V.M., Arfanakis, K., Carew, J.D., Turski, P.A., Moritz, C.H., Quigley, M.A., Meyerand, M.E., 2001. Frequencies contributing to functional connectivity in the cerebral cortex in "resting-state" data. *AJNR Am. J. Neuroradiol.* 22, 1326–1333.
- Da Cunha, C., Gomez-A. A., Blaha, C.D., 2012. The role of the basal ganglia in motivated behavior. *Rev. Neurosci.* 23 (5–6), 747–767.
- Dani, J.A., Balfour, D.J., 2011. Historical and current perspective on tobacco use and nicotine addiction. *Trends Neurosci.* 34 (7), 383–392 (Jul).
- De Lacalle, S., Iraizoz, J., Ma Gonzalo, L., 1991. Differential changes in cell size and number in topographic subdivisions of human basal nucleus in normal aging. *Neuroscience* 43, 445–456.
- Deco, G., Jirsa, V., McIntosh, A.R., Sporns, O., Kotter, R., 2009. Key role of coupling, delay, and noise in resting brain fluctuations. *Proc. Natl. Acad. Sci. U. S. A.* 106, 10302–10307.
- Delmonte, S., Gallagher, L., O'Hanlon, E., McGrath, J., Balsters, J.H., 2013. Functional and structural connectivity of frontostriatal circuitry in Autism Spectrum Disorder. *Front. Hum. Neurosci.* 7, 430 (Aug 6).
- Di Martino, A., Scheres, A., Margulies, D.S., Kelly, A.M., Uddin, L.Q., Shehzad, Z., Biswal, B., Walters, J.R., Castellanos, F.X., Milham, M.P., 2008. *Cereb. Cortex* 18 (12), 2735–2747 (Dec).
- Di Martino, A., Kelly, C., Grzadzinski, R., Zuo, X.N., Mennes, M., Mairena, M.A., Lord, C., Castellanos, F.X., Milham, M.P., 2011. Aberrant striatal functional connectivity in children with Autism. *Biol. Psychiatry* 69 (9), 847–856 (May).
- Doody, R.S., Cummings, J.L., Farlow, M.R., 2012. Reviewing the role of donepezil in the treatment of Alzheimer's disease. *Curr. Alzheimer Res.* 9 (7), 773–781 (Sep).
- Esslinger, C., Braun, U., Schirrmbeck, F., Santos, A., Meyer-Lindenberg, A., Zink, M., Kirsch, P., 2013. Activation of midbrain and ventral striatal regions implicates salience processing during a modified beads task. *PLoS One* 8 (3), e58536.
- Fair, D.A., Schlaggar, B.L., Cohen, A.L., Miezin, F.M., Dosenbach, N.U., Wenger, K.K., Fox, M.D., Snyder, A.Z., Raichle, M.E., Petersen, S.E., 2007. A method for using blocked and event-related fMRI data to study "resting state" functional connectivity. *Neuroimage* 35, 396–405.
- Ferreira, L.K., Busatto, G.F., 2013. Resting-state functional connectivity in normal brain aging. *Neurosci. Biobehav. Rev.* 37 (3), 384–400 (Mar).
- Fox, M.D., Raichle, M.E., 2007. Spontaneous fluctuations in brain activity observed with functional magnetic resonance imaging. *Nat. Rev. Neurosci.* 8, 700–711.
- Fox, M.D., Snyder, A.Z., Vincent, J.L., Corbetta, M., Van Essen, D.C., Raichle, M.E., 2005. The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proc. Natl. Acad. Sci. U. S. A.* 102, 9673–9678.
- Fox, M.D., Zhang, D., Snyder, A.Z., Raichle, M.E., 2009. The global signal and observed anticorrelated resting state brain networks. *J. Neurophysiol.* 101, 3270–3283.
- Fransson, P., 2005. Spontaneous low-frequency BOLD signal fluctuations: an fMRI investigation of the resting-state default mode of brain function hypothesis. *Hum. Brain Mapp.* 26, 15–29.
- Froemke, R.C., Carcea, I., Barker, A.J., Yuan, K., Seybold, B.A., Martins, A.R., Zaika, N., Bernstein, H., Wachs, M., Levis, P.A., Polley, D.B., Merzenich, M.M., Schreiner, C.E., 2013. Long-term modification of cortical synapses improves sensory perception. *Nat. Neurosci.* 16 (1), 79–88 (Jan).
- Fudge, J.L., Breitbart, M.A., Danish, M., Pannoni, V., 2005. Insular and gustatory inputs to the caudal ventral striatum in primates. *J. Comp. Neurol.* 490 (2), 101–118 (Sep 19).
- Garibotto, V., Tettamanti, M., Marcone, A., Florea, I., Panzacchi, A., Moresco, R., Virta, J.R., Rinne, J., Cappa, S.F., Perani, D., 2013. Cholinergic activity correlates with reserve proxies in Alzheimer's disease. *Neurobiol. Aging* 34 (11), 2694.e13–2694.e18 (Nov).
- Goard, M., Dan, Y., 2009. Basal forebrain activation enhances cortical coding of natural scenes. *Nat. Neurosci.* 12 (11), 1444–1449 (Nov).
- Gopinath, K.I., Ringe, W., Goyal, A., Carter, K., Dinse, H.R., Haley, R., Briggs, R., 2011. Striatal functional connectivity networks are modulated by fMRI resting state conditions. *Neuroimage* 54 (1), 380–388 (Jan 1).
- Grace, A.A., Floresco, S.B., Goto, Y., Lodge, D.J., 2007. Regulation of firing of dopaminergic neurons and control of goal-directed behaviors. *Trends Neurosci.* 30 (5), 220–227 (May).
- Greicius, M.D., Krasnow, B., Reiss, A.L., Menon, V., 2003. Functional connectivity in the resting brain: a network analysis of the default mode hypothesis. *Proc. Natl. Acad. Sci. U. S. A.* 100, 253–258.
- Haber, S.N., Kim, K.S., Maily, P., Calzavara, R., 2006. Reward-related cortical inputs define a large striatal region in primates that interface with associative cortical connections, providing a substrate for incentive-based learning. *J. Neurosci.* 26 (32), 8368–8376 (Aug 9).
- Hafkemeijer, A., Altmann-Schneider, I., Oleksik, A.M., van de Wiel, L., Middelkoop, H.A., van Buchem, M.A., van der Grond, J., Rombouts, S.A., 2013. Increased functional connectivity and brain atrophy in elderly with subjective memory complaints. *Brain Connect.* 3 (4), 353–362.
- Hardin, M.G., Pine, D.S., Ernst, M., 2009. The influence of context valence in the neural coding of monetary outcomes. *Neuroimage* 48 (1), 249–257 (Oct 15).
- Hedreen, J.C., Struble, R.G., Whitehouse, P.J., Price, D.L., 1984. Topography of the magnocellular basal forebrain system in human brain. *J. Neuropathol. Exp. Neurol.* 43, 1–21.
- Heimer, L., 2000. Basal forebrain in the context of schizophrenia. *Brain Res. Rev.* 31, 205–235.
- Heimer, L., de Olmos, J., Alheid, G.F., Zaborszky, L., 1991. "Perestroika" in the basal forebrain; opening the border between neurology and psychiatry. *Prog. Brain Res.* 87, 109–165.
- Henn, S., Schormann, T., Engler, K., Zilles, K., 1997. Elastische Anpassung in der digitalen Bildverarbeitung auf mehreren Auflösungsstufen mit Hilfe von Mehrgitterverfahren. In: Paulus, E., Wahl, F.M. (Eds.), *Mustererkennung*. Springer, Berlin, pp. 392–399.
- Hoffstaedter, F., Grefkes, C., Roski, C., Caspers, S., Zilles, K., Eickhoff, S.B., 2014. Age-related decrease of functional connectivity additional to gray matter atrophy in a network for movement initiation. *Brain Struct. Funct.* (Jan 8). [Epub ahead of print].
- Hoemke, L., 2006. A multigrid method for anisotropic PDEs in elastic image registration. In: Falgout, R.D. (Ed.), *Numerical Linear Algebra with Applications*. Wiley InterScience, pp. 215–229.
- Holmes, C.J., Hoge, R., Collins, L., Woods, R., Toga, A.W., Evans, A.C., 1998. Enhancement of MR images using registration for signal averaging. *J. Comput. Assist. Tomogr.* 22, 324–333.
- Honey, C.J., Kotter, R., Breakspear, M., Sporns, O., 2007. Network structure of cerebral cortex shapes functional connectivity on multiple time scales. *Proc. Natl. Acad. Sci. U. S. A.* 104, 10240–10245.
- Hu, S., Chao, H.H., Winkler, A.D., Li, C.S., 2012. The effects of age on cerebral activations: internally versus externally driven processes. *Front. Aging Neurosci.* 4, 4 (Apr 24).
- Hu, S., Chao, H.H., Zhang, S., Ide, J.S., Li, C.S., 2013. Changes in cerebral morphometry and amplitude of low-frequency fluctuations of BOLD signals during healthy aging: correlation with inhibitory control. *Brain Struct. Funct.* (Apr 4). [Epub ahead of print].
- Izhikevich, E.M., Edelman, G.M., 2008. Large-scale model of mammalian thalamocortical systems. *Proc. Natl. Acad. Sci. U. S. A.* 105, 3593–3598.
- Jenkins, G.M., Watts, D.G., 1968. *Spectral analysis and its applications*. Holden-Day, San Francisco.
- Jensen, J., McIntosh, A.R., Crawley, A.P., Mikulis, D.J., Remington, G., Kapur, S., 2003. Direct activation of the ventral striatum in anticipation of aversive stimuli. *Neuron* 40 (6), 1251–1257 (Dec 18).
- Jensen, J., Smith, A.J., Willeit, M., Crawley, A.P., Mikulis, D.J., Vitcu, I., Kapur, S., 2007. Separate brain regions code for salience vs. valence during reward prediction in humans. *Hum. Brain Mapp.* 28 (4), 294–302 (Apr).
- Kahn, I., Shohamy, D., 2013. Intrinsic connectivity between the hippocampus, nucleus accumbens, and ventral tegmental area in humans. *Hippocampus* 23 (3), 187–192 (Mar).
- Kahnt, T., Chang, L.J., Park, S.Q., Heinze, J., Haynes, J.D., 2012. Connectivity-based parcellation of the human orbitofrontal cortex. *J. Neurosci.* 32, 6240–6250.
- Kelly, A.M., Uddin, L.Q., Biswal, B.B., Castellanos, F.X., Milham, M.P., 2008. Competition between functional brain networks mediates behavioral variability. *Neuroimage* 39, 527–537.
- Kim, J.H., Lee, J.M., Jo, H.J., Kim, S.H., Lee, J.H., Kim, S.T., Seo, S.W., Cox, R.W., Na, D.L., Kim, S.I., Saad, Z.S., 2010. Defining functional SMA and pre-SMA subregions in human MFC using resting state fMRI: functional connectivity-based parcellation method. *Neuroimage* 49, 2375–2386.
- Klinkenberg, I., Blokland, A., Riedel, W.J., Sambeth, A., 2013. Cholinergic modulation of auditory processing, sensory gating and novelty detection in human participants. *Psychopharmacology (Berl)* 225 (4), 903–921 (Feb).
- Kühn, S., Gallinat, J., 2012. The neural correlates of subjective pleasantness. *Neuroimage* 61 (1), 289–294 (May 15).
- Kurosawa, R., Taoka, N., Shinohara, F., Minami, M., Kaneda, K., 2013. Cocaine exposure enhances excitatory synaptic drive to cholinergic neurons in the laterodorsal tegmental nucleus. *Eur. J. Neurosci.* (Jul 3). [Epub ahead of print].
- Laalou, F.Z., de Vasconcelos, A.P., Oberling, P., Jeltsch, H., Cassel, J.C., Pain, L., 2008. Involvement of the basal cholinergic forebrain in the mediation of general (propofol) anesthesia. *Anesthesiology* 108 (5), 888–896 (May).
- Laplante, F., Zhang, Z.W., Huppé-Gourgues, F., Dufresne, M.M., Vaucher, E., Sullivan, R.M., 2012. Cholinergic depletion in nucleus accumbens impairs mesocortical dopamine activation and cognitive function in rats. *Neuropharmacology* 63 (6), 1075–1084 (Nov).
- Leach, N.D., Nodal, F.R., Cordery, P.M., King, A.J., Bajo, V.M., 2013. Cortical cholinergic input is required for normal auditory perception and experience-dependent plasticity in adult ferrets. *J. Neurosci.* 33 (15), 6659–6671 (Apr 10).
- Leslie, F.M., Mojica, C.Y., Reynaga, D.D., 2013. Nicotinic receptors in addiction pathways. *Mol. Pharmacol.* 83 (4), 753–758 (Apr).
- Lester, D.B., Rogers, T.D., Blaha, C.D., 2010. Acetylcholine-dopamine interactions in the pathophysiology and treatment of CNS disorders. *CNS Neurosci. Ther.* 16 (3), 137–162 (Jun).
- Lin, S.C., Nicoletis, M.A., 2008. Neuronal ensemble bursting in the basal forebrain encodes salience irrespective of valence. *Neuron* 59 (1), 138–149 (Jul 10).
- Litt, A., Plassmann, H., Shiv, B., Rangel, A., 2011. Dissociating valuation and saliency signals during decision-making. *Cereb. Cortex* 21 (1), 95–102 (Jan).
- Liu, X., Hairston, J., Schrier, M., Fan, J., 2011. Common and distinct networks underlying reward valence and processing stages: a meta-analysis of functional neuroimaging studies. *Neurosci. Biobehav. Rev.* 35 (5), 1219–1236 (Apr).
- Liu, Z., Bai, L., Dai, R., Zhong, C., Wang, H., You, Y., Wei, W., Tian, J., 2012. Exploring the effective connectivity of resting state networks in mild cognitive impairment: an fMRI study combining ICA and multivariate Granger causality analysis. *Conf. Proc. IEEE Eng. Med. Biol. Soc.* 2012, 5454–5457.
- Lowe, M.J., Mock, B.J., Sorenson, J.A., 1998. Functional connectivity in single and multislice echoplanar imaging using resting-state fluctuations. *Neuroimage* 7, 119–132.
- Margulies, D.S., Kelly, A.M., Uddin, L.Q., Biswal, B.B., Castellanos, F.X., Milham, M.P., 2007. Mapping the functional connectivity of anterior cingulate cortex. *Neuroimage* 37, 579–588.
- Margulies, D.S., Vincent, J.L., Kelly, C., Lohmann, G., Uddin, L.Q., Biswal, B.B., Villringer, A., Castellanos, F.X., Milham, M.P., Petrides, M., 2009. Precuneus shares intrinsic functional architecture in humans and monkeys. *Proc. Natl. Acad. Sci. U. S. A.* 106, 20069–20074.
- Mark, G.P., Shabani, S., Dobbs, L.K., Hansen, S.T., 2011. Cholinergic modulation of mesolimbic dopamine function and reward. *Physiol. Behav.* 104 (1), 76–81 (Jul 25).
- Merker, B., 1983. Silver staining of cell bodies by means of physical development. *J. Neurosci. Methods* 9, 235–241.
- Mesulam, M.-M., Mufson, E.J., Levey, A.I., Wainer, B.H., 1983. Cholinergic innervation of cortex by the basal forebrain: cytochemistry and cortical connections of the septal area, diagonal band nuclei, nucleus basalis (substantia innominata), and hypothalamus in the Rhesus monkey. *J. Comp. Neurol.* 214, 170–197.

- Mesulam, M.M., Rosen, A.D., Mufson, E.J., 1984. Regional variations in cortical cholinergic innervation: chemoarchitectonics of acetylcholinesterase-containing fibers in the macaque brain. *Brain Res.* 311 (2), 245–258 (Oct 8).
- Mesulam, M.M., Volicser, L., Marquis, J.K., Mufson, E.J., Green, C.R., 1986. Systematic regional differences in the cholinergic innervation of the primate cerebral cortex: distribution of enzyme activities and some behavioral implications. *Ann. Neurol.* 19, 144–151.
- Miasnikov, A.A., Weinberger, N.M., 2012. Detection of an inhibitory cortical gradient underlying peak shift in learning: a neural basis for a false memory. *Neurobiol. Learn. Mem.* 98 (4), 368–379 (Nov).
- Miasnikov, A.A., Chen, J.C., Gross, N., Poytress, B.S., Weinberger, N.M., 2008. Motivationally neutral stimulation of the nucleus basalis induces specific behavioral memory. *Neurobiol. Learn. Mem.* 90 (1), 125–137 (Jul).
- Miiki, T., Kusaka, T., Yokoyama, T., Ohta, K.I., Suzuki, S., Warita, K., Jamal, M., Wang, Z.Y., Ueki, M., Liu, J.Q., Yakura, T., Tamai, M., Sumitani, K., Hosomi, N., Takeuchi, Y., 2014. Short-term ethanol exposure causes imbalanced neurotrophic factor allocation in the basal forebrain cholinergic system: a novel insight into understanding the initial processes of alcohol addiction. *J. Neural. Transm.* 121, 201–210.
- Morris, J.S., Friston, K.J., Dolan, R.J., 1997. Neural responses to salient visual stimuli. *Proc. Biol. Sci.* 264 (1382), 769–775 (May 22).
- Mufson, E.J., Ginsberg, S.D., Ikonovic, M.D., DeKosky, S.T., 2003. Human cholinergic basal forebrain: chemoanatomy and neurologic dysfunction. *J. Chem. Neuroanat.* 26 (2003), 233–242.
- Murphy, K., Birn, R.M., Handwerker, D.A., Jones, T.B., Bandettini, P.A., 2009. The impact of global signal regression on resting state correlations: are anti-correlated networks introduced? *Neuroimage* 44, 893–905.
- Nakano, K., Kayahara, T., Chiba, T., 1999. Afferent connections to the ventral striatum from the medial prefrontal cortex (area 25) and the thalamic nuclei in the macaque monkey. *Ann. N. Y. Acad. Sci.* 877, 667–670 (Jun 29).
- O'Reilly, J.X., Beckmann, C.F., Tomassini, V., Ramnani, N., Johansen-Berg, H., 2010. Distinct and overlapping functional zones in the cerebellum defined by resting state functional connectivity. *Cereb. Cortex* 20, 953–965.
- Paolone, G., Angelakos, C.C., Meyer, P.J., Robinson, T.E., Sarter, M., 2013. Cholinergic control over attention in rats prone to attribute incentive salience to reward cues. *J. Neurosci.* 33 (19), 8321–8335 (May 8).
- Pauli, W.M., O'Reilly, R.C., 2008. Attentional control of associative learning—a possible role of the central cholinergic system. *Brain Res.* 1202, 43–53 (Apr 2).
- Pohlack, S.T., Nees, F., Ruttorf, M., Schad, L.R., Flor, H., 2012. Activation of the ventral striatum during aversive contextual conditioning in humans. *Biol. Psychol.* 91 (1), 74–80 (Sep).
- Popa, D., Popescu, A.T., Pare, D., 2009. Contrasting activity profile of two distributed cortical networks as a function of attentional demands. *J. Neurosci.* 29, 1191–1201.
- Power, J.D., Barnes, K.A., Snyder, A.Z., Schlaggar, B.L., Petersen, S.E., 2012. Spurious but systematic correlations in functional connectivity MRI networks arise from subject motion. *Neuroimage* 59, 2142–2154.
- Raghandi, M.A., Simic, G., Watson, S., Stimpson, C.D., Hof, P.R., Sherwood, C.C., 2011. Comparative analysis of the nucleus basalis of Meynert among primates. *Neuroscience* 184, 1–15 (Jun 16).
- Raichle, M.E., Snyder, A.Z., 2007. A default mode of brain function: a brief history of an evolving idea. *Neuroimage* 37 (4), 1083–1090 (Oct 1).
- Raichle, M.E., MacLeod, A.M., Snyder, A.Z., Powers, W.J., Gusnard, D.A., Shulman, G.L., 2001. A default mode of brain function. *Proc. Natl. Acad. Sci. U. S. A.* 98 (2), 676–682 (Jan 16).
- Rice, M.E., Cragg, S.J., 2004. Nicotine amplifies reward-related dopamine signals in striatum. *Nat. Neurosci.* 7, 583–584.
- Robinson, S., Basso, G., Soldati, N., Sailer, U., Jovicich, J., Bruzzone, L., Kryspin-Exner, I., Bauer, H., Moser, E., 2009. A resting state network in the motor control circuit of the basal ganglia. *BMC Neurosci.* 10, 137.
- Rokem, A., Landau, A.N., Garg, D., Prinzmetal, W., Silver, M.A., 2010. Cholinergic enhancement increases the effects of voluntary attention but does not affect involuntary attention. *Neuropsychopharmacology* 35 (13), 2538–2544 (Dec).
- Rombouts, S.A., Stam, C.J., Kuijter, J.P., Scheltens, P., Barkhof, F., 2003. Identifying confounds to increase specificity during a “no task condition”. Evidence for hippocampal connectivity using fMRI. *Neuroimage* 20, 1236–1245.
- Roski, C., Caspers, S., Langner, R., Laird, A.R., Fox, P.T., Zilles, K., Amunts, K., Eickhoff, S.B., 2013. Adult age-dependent differences in resting-state connectivity within and between visual-attention and sensorimotor networks. *Front. Aging Neurosci.* 5, 67 (Oct 29).
- Rothkirch, M., Schmack, K., Schlagenhauf, F., Sterzer, P., 2012. Implicit motivational value and salience are processed in distinct areas of orbitofrontal cortex. *Neuroimage* 62 (3), 1717–1725 (Sep).
- Sarter, M., Gehring, W.J., Kozak, R., 2006. More attention must be paid: the neurobiology of attentional effort. *Brain Res. Rev.* 51 (2), 145–160 (Aug).
- Scholvinck, M.L., Maier, A., Ye, F.Q., Duyn, J.H., Leopold, D.A., 2010. Neural basis of global resting-state fMRI activity. *Proc. Natl. Acad. Sci. U. S. A.* 107, 10238–10243.
- Schormann, T., Zilles, K., 1998. Three-dimensional linear and nonlinear transformations: An integration of light microscopical and MRI data. *Hum. Brain Map.* 6, 339–347.
- Schwarz, A.J., McGonigle, J., 2011. Negative edges and soft thresholding in complex network analysis of resting state functional connectivity data. *Neuroimage* 55, 1132–1146.
- Selden, N.R., Gitelman, D.R., Salamon-Murayama, N., Parrish, T.B., Mesulam, M.M., 1998. Trajectories of cholinergic pathways within the cerebral hemispheres of the human brain. *Brain* 121 (Pt 12), 2249–2257.
- Smyser, C.D., Inder, T.E., Shimony, J.S., Hill, J.E., Degnan, A.J., Snyder, A.Z., Neil, J.J., 2010. Longitudinal analysis of neural network development in preterm infants. *Cereb. Cortex* 20, 2852–2862.
- Sofuoglu, M., Mooney, M., 2009. Cholinergic functioning in stimulant addiction: implications for medications development. *CNS Drugs* 23 (11), 939–952 (Nov).
- Spicer, J., Galvan, A., Hare, T.A., Voss, H., Glover, G., Casey, B., 2007. Sensitivity of the nucleus accumbens to violations in expectation of reward. *Neuroimage* 34 (1), 455–461 (Jan 1).
- Taniwaki, T., Okayama, A., Yoshiura, T., Togao, O., Nakamura, Y., Yamasaki, T., Ogata, K., Shigetou, H., Ohyagi, Y., Kira, J., Tobimatsu, S., 2007. Age-related alterations of the functional interactions within the basal ganglia and cerebellar motor loops in vivo. *Neuroimage* 36 (4), 126 (Jul 15).
- Thiel, C.M., Bentley, P., Dolan, R.J., 2002. Effects of cholinergic enhancement on conditioning-related responses in human auditory cortex. *Eur. J. Neurosci.* 16 (11), 2199–2206 (Dec).
- Threlfell, S., Cragg, S.J., 2011. Dopamine signaling in dorsal versus ventral striatum: the dynamic role of cholinergic interneurons. *Front. Syst. Neurosci.* 5, 11 (Mar 3).
- Tomasi, D., Volkow, N.D., 2014. Functional connectivity of substantia nigra and ventral tegmental area: maturation during adolescence and effects of ADHD. *Cereb. Cortex* 24, 935–944.
- Uddin, L.Q., Kelly, A.M., Biswal, B.B., Xavier Castellanos, F., Milham, M.P., 2009. Functional connectivity of default mode network components: correlation, anticorrelation, and causality. *Hum. Brain Mapp.* 30, 625–637.
- Van Dijk, K.R., Sabuncu, M.R., Buckner, R.L., 2012. The influence of head motion on intrinsic functional connectivity MRI. *Neuroimage* 59, 431–438.
- Veldhuizen, M.G., Douglas, D., Aschenbrenner, K., Gitelman, D.R., Small, D.M., 2011. The anterior insular cortex represents breaches of taste identity expectation. *J. Neurosci.* 31 (41), 14735–14744 (Oct 12).
- Vogels, O.J.M., Broere, C.A.J., Ter Laak, H.J., Ten Donkelaar, H.J., Nieuwenhuys, R., Schulte, B.P.M., 1990. Cell loss and shrinkage in the nucleus basalis Meynert complex in Alzheimer's disease. *Neurobiol. Aging* 11, 3–13.
- Wallace, T.L., Bertrand, D., 2013. Importance of the nicotinic acetylcholine receptor system in the prefrontal cortex. *Biochem. Pharmacol.* 85 (12), 1713–1720 (Jun 15).
- Weissenbacher, A., Kasess, C., Gerstl, F., Lanzenberger, R., Moser, E., Windischberger, C., 2009. Correlations and anticorrelations in resting-state functional connectivity MRI: a quantitative comparison of preprocessing strategies. *Neuroimage* 47, 1408–1416.
- Wenk, G.L., 1997. The nucleus basalis magnocellularis cholinergic system: one hundred years of progress. *Neurobiol. Learn. Mem.* 67 (2), 85–95 (Mar).
- Williams, M.J., Adinoff, B., 2008. The role of acetylcholine in cocaine addiction. *Neuropsychopharmacology* 33 (8), 1779–1797 (Jul).
- Wilson, F.A., Ma, Y.Y., 2004. Reinforcement-related neurons in the primate basal forebrain respond to the learned significance of task events rather than to the hedonic attributes of reward. *Cogn. Brain Res.* 19 (1), 74–81.
- Wilson, F.A., Rolls, E.T., 1990. Neuronal responses related to reinforcement in the primate basal forebrain. *Brain Res.* 509 (2), 213–231.
- Ystad, M., Eichele, T., Lundervold, A.J., Lundervold, A., 2010. Subcortical functional connectivity and verbal episodic memory in healthy elderly—a resting state fMRI study. *Neuroimage* 52 (1), 379–388 (Aug 1).
- Zaborszky, L., Cullinan, W.E., 1992. Projections from the nucleus accumbens to cholinergic neurons of the ventral pallidum: a correlated light and electron microscopic double immunolabeling study in rat. *Brain Res.* 570, 92–101.
- Zaborszky, L., Hoemke, L., Mohlberg, H., Schleicher, A., Amunts, K., Zilles, K., 2008. Stereotaxic probabilistic maps of the magnocellular cell groups in human basal forebrain. *Neuroimage* 42, 1127–1141.
- Zaborszky, L., Van den Pol, A., Gyengesi, E., 2012. The basal forebrain cholinergic projection system in mice. In: Watson, C., Paxinos, G., Puelles, L. (Eds.), *The Mouse Nervous System*, 1st ed. Elsevier, Amsterdam, pp. 684–718.
- Zaborszky, L., Csordas, A., Mosca, K., Kim, J., Gielow, M.R., Vadasz, C., Nadasdy, Z., 2014. Neurons in the basal forebrain project to the cortex in a complex topographic organization that reflects corticocortical connectivity patterns: an experimental study based on retrograde tracing and 3D reconstruction. *Cereb. Cortex* (in press, 2013 Aug 19. [Epub ahead of print]).
- Zhang, S., Li, C.S., 2012a. Functional connectivity mapping of the human precuneus by resting state fMRI. *Neuroimage* 59, 3548–3562.
- Zhang, S., Li, C.S., 2012b. Functional networks for cognitive control in a stop signal task: independent component analysis. *Hum. Brain Mapp.* 33, 89–104.
- Zhang, S., Li, C.-S., 2014. Functional clustering of the human inferior parietal lobule by whole-brain connectivity mapping of resting-state functional magnetic resonance imaging signals. *Brain Connect.* 4 (1), 53–69.
- Zhang, D., Snyder, A.Z., Fox, M.D., Sansbury, M.W., Shimony, J.S., Raichle, M.E., 2008. Intrinsic functional relations between human cerebral cortex and thalamus. *J. Neurophysiol.* 100, 1740–1748.
- Zhang, D., Snyder, A.Z., Shimony, J.S., Fox, M.D., Raichle, M.E., 2010. Noninvasive functional and structural connectivity mapping of the human thalamocortical system. *Cereb. Cortex* 20, 1187–1194.
- Zhang, S., Ide, J.S., Li, C.S., 2012. Resting-state functional connectivity of the medial superior frontal cortex. *Cereb. Cortex* 22, 99–111.
- Zink, C.F., Pagnoni, G., Martin, M.E., Dhamala, M., Berns, G.S., 2003. Human striatal response to salient nonrewarding stimuli. *J. Neurosci.* 23 (22), 8092–8097 (Sep 3).
- Zink, C.F., Pagnoni, G., Chappelow, J., Martin-Skurski, M., Berns, G.S., 2006. Human striatal activation reflects degree of stimulus saliency. *Neuroimage* 29 (3), 977–983 (Feb 1).
- Zuo, X.N., Kelly, C., Adelstein, J.S., Klein, D.F., Castellanos, F.X., Milham, M.P., 2010. Reliable intrinsic connectivity networks: test-retest evaluation using ICA and dual regression approach. *Neuroimage* 49, 2163–2177.